

FRONT PANEL

Kill-All Pellets

Reg. No: **L7162** Act/Wet No 36 of /van 1947

**RAT AND MOUSE PELLETS
ROT EN MUIS KORRELS**

For the control of the Norway rat (*Rattus norvegicus*),
roof rat (*Rattus rattus*) and the house mouse (*Mus musculus*).
Vir die beheer van Norweegse rot (*Rattus norvegicus*),
dakrot (*Rattus rattus*) en die huismuis (*Mus musculus*).

Active ingredient:

Aktiewe bestanddeel:

Brodifacoum (anticoagulant)....0,02 g/kg

Brodifakum (antikoagulant)....0,02 g/kg

REGISTRATION HOLDER / REGISTRASIEHOUER:

Almond Agro Chemicals (Pty) Ltd / (Edms) Bpk

Reg. No. 1997/006442/07

P.O.Box / Posbus 188,

Van Riebeeck Park, 1629

Lot No.:

Expiry date / Verval datum:

500 g

HARMFUL / SKADELIK

[Yellow colour band]

SIDE PANEL

WARNINGS: Handle with care. Keep out of reach of children, uninformed persons and animals. Store away from food and feedstuffs. Poisonous if swallowed and upon prolonged contact with the skin. Toxic to fish, birds and animals. In case of poisoning, call a doctor and make this label available to him.

PRECAUTIONS: Avoid contact with skin. Wear gloves when handling product. Avoid contamination of any food, food utensils and drinking water. Place bait in locations out of reach of children and animals. Wash hands and exposed skin after handling. Destroy empty container and do not re-use for any other purpose. Bury dead rodents and any bait not consumed at end of treatment.

SYMPTOMS OF POISONING: Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints, and hematomata. Blood in urine and faeces, and abdominal pain may occur.

FIRST AID TREATMENT: Remove patient from source of poisoning and keep him quiet and reassured. Take patient IMMEDIATELY to the nearest physician. If person is unconscious, do not give anything by mouth and do not induce vomiting.

NOTE TO PHYSICIAN: Administer 5 mg/kg Vit. K₁ intravenously on the first day, followed by another administration of the same dose if the Quick time does not normalize a few hours after the first injection. Follow up with 5 mg/kg/day orally on the second and the following days. The duration of the treatment is generally between 10 and 15 days. In severe cases give small transfusions of fresh blood.

DIRECTIONS FOR USE: Use only as directed.

For use in the home, on the farm and industrial premises. **Kill-All Pellets** can kill rats and mice after a single feed. Death of rodents will take 4 – 12 days.

- 1. Norway and Roof Rats:** Place **Kill-All Pellets** in a covered bait station to prevent access by children and domestic animals. Place 50 – 75 g **Kill-All Pellets** per bait station every 5 – 10 m along walls or ratruns, in rat burrows and in sheltered areas where rats are known to live or feed. Bait points should be inspected and replenished every 7 days until feeding ceases.
- 2. Mice:** Place 15 – 20 g **Kill-All Pellets** per bait station every 2 – 4 m along walls or runways, in nests and in sheltered areas where mice are known to live or feed. The single feed action of **Kill-All Pellets** is particularly effective against mice with their small appetites and sporadic feeding behaviour.